

# Repeat Selective Laser Trabeculoplasty

Bryan Kun Hong, BS,\* Jeffrey C. Winer, MD,\* James F. Martone, MD, MPH,†  
Martin Wand, MD,‡§ Bruce Altman, MD,|| and Bruce Shields, MD\*¶

**Purpose:** To evaluate the efficacy of repeat 360-degree selective laser trabeculoplasty (SLT) in glaucoma patients with prior successful 360-degree SLT.

**Design:** Retrospective chart review.

**Methods:** Forty-four eyes of 35 patients,  $\geq 18$  years of age, with open-angle glaucoma (primary open-angle, pseudoexfoliation, or pigmentary glaucoma), uncontrolled on maximum tolerable medical therapy, underwent an initial 360-degree SLT (SLT1), which was successful for  $\geq 6$  months, but eventually lost efficacy and was followed by a repeat 360-degree SLT (SLT2). Patients with prior argon laser trabeculoplasty or other glaucoma surgery, before or during the study period, were excluded. Intraocular pressure (IOP) measurements were recorded before each procedure and 1 to 4 weeks, 1 to 3 months, and 5 to 8 months posttreatment and 15 to 21 weeks after the initial SLT.

**Results:** Reduction in IOP after SLT1 and SLT2 was significantly less with repeat treatment at 1 to 3 months, with average decreases of  $-5.0$  and  $-2.9$  mm Hg, respectively ( $P = 0.01$ ), but there were no statistically significant differences between treatments at the other equivalent time points. Using a definition of "success" as  $\geq 20\%$  peak IOP reduction, success rates for SLT1 and SLT2 were not significantly different. There was also no significant difference in eyes that received SLT2 6 to 12 months after SLT1 compared with those that received SLT2 12 months or more after SLT1.

**Conclusions:** Our findings suggest that repeat 360-degree SLT may be safe and effective after an initially successful 360-degree SLT has failed. These results may be achieved as early as 6 months after the first treatment.

**Key Words:** selective laser trabeculoplasty, SLT, repeat SLT

(*J Glaucoma* 2009;18:180–183)

Selective laser trabeculoplasty (SLT) is becoming an increasingly popular alternative to argon laser trabeculoplasty (ALT) in the treatment of open-angle forms of glaucoma, primarily owing to the lack of collateral thermal damage and trabecular meshwork scarring with SLT as compared with ALT.<sup>1</sup> However, prospective clinical trials

have shown no significant difference between the 2 treatments with regard to safety and efficacy upon initial treatment.<sup>2</sup> A pressing question is whether repeating SLT, after an initially successful SLT has lost its efficacy, is associated with further clinically significant intraocular pressure (IOP) reduction, which has not been found to be the case with ALT.<sup>3</sup>

We report the results of a retrospective study in which a repeat 360-degree SLT was performed 6 months or more after an initially successful 360-degree SLT.

## METHODS

This is a retrospective chart review of patients treated in the offices of 3 of the authors (B.A., J.M., and M.W.). Inclusion criteria were open-angle glaucoma patients (primary open-angle, pseudoexfoliation, or pigmentary glaucoma),  $\geq 18$  years of age, who were uncontrolled on maximum tolerable medical therapy and underwent an initial 360-degree SLT (SLT1), which was successful for 6 months or more, but eventually lost efficacy and was followed by a repeat 360-degree SLT (SLT2). Exclusion criteria were prior ALT or other glaucoma surgery before or during the study period. The study was approved by the Human Investigation Committee of Yale University School of Medicine.

Data recorded for each patient included age, sex, type of glaucoma, glaucoma medications just before and during the study, and the SLT protocol (number of spots and laser power setting). Baseline IOP for SLT1 was taken as the average of the last 3 readings before the treatment. The baseline IOP for SLT2 was the measurement on the day of the repeat treatment.

The SLT protocol consisted of pretreating with apraclonidine and applying laser energy to all 4 quadrants of the trabecular meshwork. All eyes were treated with bromfenac or diclofenac b.i.d. for 4 to 7 days post-SLT. Follow-up IOP measurements were obtained at 1 to 4 weeks, 1 to 3 months, and 5 to 8 months for both SLT1 and SLT2.

To evaluate the influence of time after SLT1 before the repeat procedure, the patients were divided into those who received SLT2 6 to 12 months after SLT1 (early repeats) and those in which the repeat was performed more than 12 months after SLT1 (late repeats). In addition to the 3 time points, the early and late repeat groups were compared at 15 to 21 months. The minimum follow-up time for all patients was 6 months after both SLT1 and SLT2.

The main outcome variables analyzed were the IOP-lowering effect of SLT2 compared with SLT1 at each time point and the IOP-lowering effect of SLT2 in the early repeat group as compared with that of the late repeat group. SLT1 and SLT2 were compared, using a definition of "success" as  $\geq 20\%$  IOP reduction in relation to

Received for publication January 9, 2008; accepted May 3, 2008.

From the \*Yale University School of Medicine; †Yale Eye Center, New Haven; ‡The Eye Center, A Medical and Surgical Group, PC, Hamden; §Department of Ophthalmology, University of Connecticut School of Medicine; ¶Consulting Ophthalmologists, Farmington; and ||Danbury Eye Physicians and Surgeons, PC, Danbury, CT.

Financial Disclosures: None.

Supported by National Institutes of Health-NCRR CTSA-T32 Medical Student Research Fellowship and Yale University School of Medicine Medical Student Research Fellowship.

Reprints: Bruce Shields, MD, Yale Eye Center, 40 Temple Street, New Haven, CT 06510 (e-mail: bruce.shields@yale.edu).

Copyright © 2009 by Lippincott Williams & Wilkins

respective pretreatment baselines. IOP increases above the aforementioned threshold of success were seen as loss of efficacy. To compare the efficacy from an equivalent time point relative to the respective procedures, the baseline for SLT2 is taken to be the IOP measurement directly before the second surgery. This provides a null hypothesis that the previous surgery does not affect the efficacy of repeat surgery. Baseline characteristic distributions were also compared between the early repeat and late repeat groups.

Student *t* test was used to evaluate statistical significance in continuous variables in which there was a single measurement. In cases where the SLT1 baseline measurements were compared, because there were 3 measurements, *P* values were calculated using repeat measures analysis using analysis of variance.  $\chi^2$  testing was used to calculate significance in categorical variables. All of these calculations were performed using Data Desk version 6.1 by Data Description Inc.

As we have 9 patients in whom both eyes were included, we first analyzed right and left eyes separately to avoid confounding of variables and pooled the data upon finding that the results were not significantly different.

### RESULTS

Forty-four eyes of 35 patients met the inclusion-exclusion criteria and were included in the study. Patient demographics are shown in Table 1. Table 2 shows the mean IOP and mean IOP changes for SLT1 and SLT2 at the 3 time points for all 44 eyes. The only statistically significant difference was the mean change at 1 to 3 months, which was  $-5.0$  mm Hg for SLT1 and  $-2.9$  mm Hg for SLT2 ( $P = 0.01$ ). Table 3 shows the percentage of eyes that qualified for the definition of success ( $\geq 20\%$  IOP reduction) at each time point. At the 5 to 8-month time point, the success rate for SLT1 was 7% higher than that for SLT2, but the difference did not reach statistical significance. Eight patients (18.2%) had IOP greater than their SLT2 baselines 1 to 4 weeks after second treatment; of these patients, however, only 1 eye (2.3%) continued to have elevated IOP through the remainder of follow-up. With the exception of 1 patient, in whom brimonidine was discontinued owing to dizziness, there were no changes in the glaucoma medication regimens throughout the study.

Mild anterior uveitis was documented in 2 eyes post-SLT1 and an IOP spike was noted in 1 (2.3%) eye after both SLT1 and SLT2. Conjunctival hyperemia was noted in 26 (59%) eyes post-SLT1 and 28 (64%) eyes post-SLT2. Posttreatment complications were not significantly different between the 2 surgeries.

**TABLE 2.** Intraocular Pressures at Equivalent Time Points After Initial Application of Selective Laser Trabeculoplasty (SLT1) and After Repeat Application (SLT2) in the Treatment of Open-angle Glaucoma and Change From Preprocedure Baseline IOP (Pre-SLT2 IOP for SLT2 Measurements) (n = 44)

		SLT1	SLT2	<i>P</i>
IOP	Baseline	20.1 (18.8, 21.3)	19.5 (17.7, 21.3)	0.49
	1-4 wk	16.9 (15.7, 18.2)	16.9 (15.1, 18.6)	0.94
	1-3 mo	15.0 (14.2, 15.8)	16.6 (15.2, 18.0)	0.04
IOP change	5-8 mo	16.2 (14.9, 17.4)	16.7 (14.9, 18.4)	0.60
	1-4 wk	-3.1 (-4.6, -1.7)	-2.6 (-4.1, -1.1)	0.61
	1-3 mo	-5.0 (-6.2, -3.9)	-2.9 (-4.0, -1.8)	0.002
	5-8 mo	-4.0 (-5.3, -2.7)	-2.9 (-4.2, -1.5)	0.16

IOP indicates intraocular pressure; SLT, selective laser trabeculoplasty.

The demographics for the early repeat and late repeat groups are also shown in Table 1. The only significant difference was the distribution of right and left eyes. The baseline mean IOPs, mean IOPs, and mean change in IOPs at the 4 time points for the 2 groups are shown in Table 4. Although there was a statistically significant difference in baseline values, being higher in the early repeat group, the only significant difference in the posttreatment time periods was a lower mean IOP in the late repeat group at 5 to 8 months.

### DISCUSSION

Damji et al<sup>2</sup> studied SLT after ALT and reported beneficial IOP reduction with SLT in eyes that had been unresponsive to ALT. Two groups (Mequio MJ et al IOVS 2007; 48: ARVO E-Abstract 3972 and Franco TC et al IOVS 2007; 48: ARVO E-Abstract 3981) recently described their experience with repeat SLT after prior SLT, although the initial SLTs were not all 360-degree and the repeat procedures were performed in some quadrants that had been previously treated and some that had not. To our knowledge, there has been no previous report in which a first 360-degree SLT, which was initially successful, but eventually lost efficacy, was followed by a repeat 360-degree SLT.

There is much written in the literature as to the viability of repeat ALT. Early studies suggested that repeat 360-degree ALT in previously failed 360-degree ALT yielded encouraging results, with a 100% success rate at 3 months in 1 study<sup>4</sup> and up to 53% of retreated eyes achieving an IOP reduction at 12 weeks in another.<sup>5</sup>

**TABLE 1.** Initial Demographics: Mean (95% CI) or Number (%) of All Glaucomatous Eyes and 2 Treatment Groups (Selective Laser Trabeculoplasty Repeated at 6 to 12 mo and  $\geq 12$  mo)

	Total	6-12 mo	$\geq 12$ mo	<i>P</i>
Sex (M:F)	15:29 (34:66)	7:13 (35:65)	8:16 (33:67)	0.91
Age	73.2 (69.7, 76.9)	73.6 (67.9, 79.3)	72.99 (68.3, 77.7)	0.45
Type of Glaucoma (POAG:PXG:PG)	35:8:1 (80:18:12)	17:2:1 (85:10:5)	18:6:0 (75:25:0)	0.26
Eye (OD:OS)*	25:19 (57:43)	14:6 (70:30)	11:13 (46:54)	0.006
No. Spots	102.1 (98.1, 106.0)	102.4 (96.3, 108.5)	101.8(96.5, 107.1)	0.88
Power setting (mJ)	88.1 (84.7, 91.5)	87.8 (82.3, 93.3)	88.4 (84.2, 92.6)	0.87

\*Right eye:left eye.

CI indicates confidence interval; F, female; M, male; PG, pigmentary glaucoma; POAG, primary open-angle glaucoma; PXG, pseudoexfoliative glaucoma.

**TABLE 3.** Percentage of Glaucomatous Eyes That Reach  $\geq 20\%$  Decrease in Intraocular Pressure After Initial Application of Selective Laser Trabeculoplasty (SLT1) and After Repeat Application (SLT2)

	1-4 wk	1-3 mo	5-8 mo
SLT 1	29.5% (n = 13)	54.5% (n = 24)	50% (n = 22)
SLT 2	40.9% (n = 18)	36.3% (n = 16)	43.2% (n = 19)
	<i>P</i> = 0.26	<i>P</i> = 0.09	<i>P</i> = 0.52

SLT indicates selective laser trabeculoplasty.

However, these numbers prove to be a bit misleading in that the former figure considers treatment to be successful if IOP is maintained below an absolute pressure of 21 mm Hg for all eyes and the latter figure takes even the smallest IOP reductions into account.

In a study regarding repeat ALT in eyes that had a positive prolonged response to initial 180-degree and 360-degree ALT, Jorizzo et al<sup>6</sup> showed that 8 of 11 eyes had sustained reduction of IOP at >1-year follow-up. Although this too is encouraging, we cannot be sure whether the IOP-lowering effect came from retreatment of previously treated regions or whether the apparent effect is skewed by the treatment of previously untreated regions (ie, 180-degree ALT1). It is appearing more and more likely that the therapeutic benefits of repeat ALT are limited<sup>7</sup> and that, although fairly safe, the likelihood of successful IOP reduction decreases significantly with each repeat ALT.<sup>3</sup>

On the other hand, SLT has been shown to be better tolerated, with less discomfort and postoperative anterior chamber inflammation<sup>8</sup> and significantly less thermal damage to the trabecular meshwork as compared with ALT.<sup>9</sup>

We found that a repeat 360-degree SLT provided additional IOP reduction, which was not as marked as that with the first treatment, although the difference between the 2 was not statistically significant except at the 1 to 3-month time period. We also found that only 1 eye (2.3%) continued to have elevated IOP through the remainder of post-SLT2 follow-up.

We chose to be consistent with Damji et al<sup>2</sup> in defining success as being a  $\geq 20\%$  reduction of pretreatment IOP. Although our follow-up was not necessarily  $\geq 1$  year after each SLT application, we felt that this definition of success

would not only make our results comparable with those of previous studies, but also account for those IOP reductions closely approaching the average rather than only those IOP reductions greater than or equal to the average. With this definition of success, there was again no significant difference between the initial and repeat treatments. We also found that the time since the initial SLT did not influence the outcome of the repeat procedure.

Even though patients who had repeat SLT within 6 to 12 months of the initial treatment had a significantly higher baseline IOP than those whose repeat was after 12 months, the only significant difference between these 2 groups was a lower mean pressure in the late repeat group at the 5 to 8-month time period.

Our study is limited by its retrospective design. The decision to repeat SLT was made by the treating physician on the basis of the overall status of each patient. The sample size is also fairly small, so larger, prospective studies are needed to confirm our findings. In addition, because 360-degree repeat SLT has been so rare to date, it was necessary to accept using multiple eyes from the same subjects to gain proper power for this retrospective trial. Although not ideal, there were neither differences between the 2 eyes of those repeated subjects, nor between the demographics of them and those subjects who had repeat SLT in only 1 eye. Another measure that allowed for increased power was the fact that 3 measurements were taken before SLT1. Because the IOP before SLT1 was in steady state, 3 measurements were taken to better calculate the baseline; however, between the first and second surgeries, at the time of SLT2 baseline, the IOP was in a state of flux, so repeat measure would have been misleading. The repeated measures were not significantly different from each other as judged by analysis of variance with patient ID and order as factors (*P* = 0.54).

Nevertheless, our preliminary findings are encouraging, suggesting that 360-degree SLT may be repeated after an initially successful 360-degree SLT has failed, with IOP reduction that suggests only slightly diminishing returns. Our findings also suggest that these results may be achieved as early as 6 months after the first treatment. If prospective, randomized trials also show that SLT can be repeated in a safe and effective manner, it may be possible to claim an advantage for the use of SLT over ALT, and the former may assume an increasingly important role in our management of glaucoma.

**TABLE 4.** Comparison of Intraocular Pressure Reduction With Repeat Selective Laser Trabeculoplasty (SLT2) 6-12 mo and  $\geq 12$  mo After SLT1 in the Treatment of Open-angle Glaucoma (95% CI\*)

		6-12 mo SLT2	$\geq 12$ mo SLT2	<i>P</i>
IOP	Preoperative	21.7 (19.4, 24.0)	18.7 (17.7, 19.76)	0.01
	1-4 wk	17.4 (15.7, 19.0)	16.6 (14.8, 18.4)	0.55
	1-3 mo	15.5 (14.3, 16.6)	14.7 (13.6, 15.7)	0.33
	5-8 mo	17.5 (15.6, 19.4)	15.0 (13.6, 16.5)	0.02
	15-21 mo	18.1 (14.7, 21.4)	17.0 (15.7, 18.4)	0.56
	IOP change	1-4 wk	-4.3 (-6.7, -2.0)	-2.1 (-3.9, -0.4)
1-3 mo		-6.2 (-8.1, -4.3)	-4.1 (-5.5, -2.7)	0.07
5-8 mo		-4.2 (-6.3, -2.1)	-3.9 (-5.6, -2.2)	0.41
15-21 mo		-3.6 (-7.0, -0.2)	-1.7 (-3.3, -0.1)	0.29

\*Confidence interval.

IOP indicates intraocular pressure; SLT, selective laser trabeculoplasty.

## REFERENCES

1. Latina M, Park C. Selective targeting of trabecular meshwork cells: in vitro studies of pulsed and CW laser interactions. *Exp Eye Res.* 1995;60:359–372.
2. Damji KF, Shah KC, Rock WJ, et al. Selective laser trabeculoplasty vs. argon laser trabeculoplasty: a prospective randomized clinical trial. *Br J Ophthalmol.* 1999;83:718–722.
3. Richter CU, Shingleton BJ, Bellows AR, et al. Retreatment with argon laser trabeculoplasty. *Ophthalmology.* 1987;94:1085–1089.
4. Weber PA, Burton GD, Epitropoulos AT. Laser trabeculoplasty retreatment. *Ophthalmic Surg.* 1989;20:702–706.
5. Starita RJ, Fellman RL, Spaeth GL, et al. The effect of repeating full-circumference argon laser trabeculoplasty. *Ophthalmic Surg.* 1984;15:41–43.
6. Jorizzo PA, Samples JR, Van Buskirk EM. The effect of repeat argon laser trabeculoplasty. *Am J Ophthalmol.* 1988;106:682–685.
7. Fink AI, Jordan AJ, Lao PN, et al. Therapeutic limitations of argon laser trabeculoplasty. *Br J Ophthalmol.* 1988;72:263–269.
8. Martinez-de-la-Casa JM, Garcia-Feijoo J, Castillo A, et al. Selective vs argon laser trabeculoplasty: hypotensive efficacy, anterior chamber inflammation, and postoperative pain. *Eye.* 2004;18:498–502.
9. Kramer TR, Noecker RJ. Comparison of the morphologic changes after selective laser trabeculoplasty and argon laser trabeculoplasty in human eye bank eyes. *Ophthalmology.* 2001;108:773–779.